

Dephosphorylation of compounds **70** and **74**, **75** and **76** was accomplished by treating 10 µl of the crude stock solutions (ranging in concentration from approximately 0.5 to 2 mM) with 2 units of CIAP in 100 µl of CIAP buffer (Promega) at 37°C for 1 hour. The reactions were then heated to 75°C for 15 min. in order to inactivate the CIAP. For clarity, dephosphorylated compounds are designated 'dp'. For example, after dephosphorylation, substrate **70** becomes **70dp**.

To prepare samples for IEF experiments, the concentration of the stock solutions of substrate and dephosphorylated product were adjusted to a uniform absorbance of  $8.5 \times 10^{-3}$  at 532 nm by dilution with water. Two microliters of each sample were analyzed by IEF using a PhastSystem electrophoresis unit (Pharmacia) and PhastGel IEF 3-9 media (Pharmacia) according to the manufacturer's protocol. Separation was performed at 15°C with the following program: pre-run; 2,000 V, 2.5 mA, 3.5 W, 75 Vh; load; 200 V, 2.5 mA, 3.5 W, 15 Vh; run; 2,000 V; 2.5 mA; 3.5 W, 130 Vh. After separation, samples were visualized by using the FMBIO Image Analyzer (Hitachi) fitted with a 585 nm filter. The resulting imager scan is shown in Figure 59.

Figure 59 shows results of IEF separation of substrates **70**, **74**, **75** and **76** and their dephosphorylated products. The arrow labeled "Sample Loading Position" indicates a loading line, the '+' sign shows the position of the positive electrode and the '-' sign indicates the position of the negative electrode.

The results shown in Figure 59 demonstrate that substrates **70**, **74**, **75** and **76** migrated toward the positive electrode, while the dephosphorylated products **70dp**, **74dp**, **75dp** and **76dp** migrated toward negative electrode. The observed differences in mobility direction was in accord with predicted net charge of the substrates (minus one) and the products (plus one). Small perturbations in the mobilities of the phosphorylated compounds indicate that the overall pIs vary. This was also true for the dephosphorylated compounds. The presence of the cytosine in **76dp**, for instance, moved this compound further toward the negative electrode which was indicative of a higher overall pI relative to the other dephosphorylated compounds. It is important to note that additional positive charges can be obtained by using a combination of natural

amino modified bases (70dp and 74dp) along with uncharged methylphosphonate bridges (products 75dp and 76dp).

The results shown above demonstrate that the removal of a single phosphate group can flip the net charge of an oligonucleotide to cause reversal in an electric field, allowing easy separation of products, and that the precise base composition of the oligonucleotides affect absolute mobility but not the charge-flipping effect.

### EXAMPLE 23

#### Detection Of Specific Cleavage Products In The Invader-Directed Cleavage Reaction By Charge Reversal

In this example the ability to isolate products generated in the invader-directed cleavage assay from all other nucleic acids present in the reaction cocktail was demonstrated using charge reversal. This experiment utilized the following Cy3-labeled oligonucleotide: 5'-Cy3-AminoT-AminoT-CTTTTCACCAGCGAGACGGG-3' (SEQ ID NO:61; termed "oligo 61"). Oligo 61 was designed to release upon cleavage a net positively charged labeled product. To test whether or not a net positively charged 5'-end labeled product would be recognized by the Cleavase® enzymes in the invader-directed cleavage assay format, probe oligo 61 (SEQ ID NO:61) and invading oligonucleotide 67 (SEQ ID NO:62) were chemically synthesized on a DNA synthesizer (ABI 391) using standard phosphoramidite chemistries and reagents obtained from Glen Research (Sterling, VA).

Each assay reaction comprised 100 fmoles of M13mp18 single stranded DNA, 10 pmoles each of the probe (SEQ ID NO:61) and invader (SEQ ID NO:62) oligonucleotides, and 20 units of Cleavase® A/G in a 10 µl solution of 10 mM MOPS, pH 7.4 with 100 mM KCl. Samples were overlaid with mineral oil to prevent evaporation. The samples were brought to either 50°C, 55°C, 60°C, or 65°C and cleavage was initiated by the addition of 1 µl of 40 mM MnCl<sub>2</sub>. Reactions were allowed to proceed for 25 minutes and then were terminated by the addition of 10 µl of 95% formamide containing 20 mM EDTA and 0.02% methyl violet. The negative

control experiment lacked the target M13mp18 and was run at 60°C. Five microliters of each reaction were loaded into separate wells of a 20% denaturing polyacrylamide gel (cross-linked 29:1) with 8 M urea in a buffer containing 45 mM Tris-Borate (pH 8.3) and 1.4 mM EDTA. An electric field of 20 watts was applied for 30 minutes, with the electrodes oriented as indicated in Figure 60B (*i.e.*, in reverse orientation). The products of these reactions were visualized using the FMBIO fluorescence imager and the resulting imager scan is shown in Figure 60B.

Figure 60A provides a schematic illustration showing an alignment of the invader (SEQ ID NO:61) and probe (SEQ ID NO:62) along the target M13mp18 DNA; only 53 bases of the M13mp18 sequence is shown (SEQ ID NO:63). The sequence of the invader oligonucleotide is displayed under the M13mp18 target and an arrow is used above the M13mp18 sequence to indicate the position of the invader relative to the probe and target. As shown in Figure 60A, the invader and probe oligonucleotides share a 2 base region of overlap.

In Figure 60B, lanes 1-6 contain reactions performed at 50°C, 55°C, 60°C, and 65°C, respectively; lane 5 contained the control reaction (lacking target). In Figure 60B, the products of cleavage are seen as dark bands in the upper half of the panel; the faint lower band seen appears in proportion to the amount of primary product produced and, while not limiting the invention to a particular mechanism, may represent cleavage one nucleotide into the duplex. The uncleaved probe does not enter the gel and is thus not visible. The control lane showed no detectable signal over background (lane 5). As expected in an invasive cleavage reaction, the rate of accumulation of specific cleavage product was temperature-dependent. Using these particular oligonucleotides and target, the fastest rate of accumulation of product was observed at 55°C (lane 2) and very little product observed at 65°C (lane 4).

When incubated for extended periods at high temperature, DNA probes can break non-specifically (*i.e.*, suffer thermal degradation) and the resulting fragments contribute an interfering background to the analysis. The products of such thermal breakdown are distributed from single-nucleotides up to the full length probe. In this experiment, the ability of charge based separation of cleavage products (*i.e.*, charge